wellcare

health net

Use of Imaging Studies for Low Back Pain

Wellcare by Health Net (Health Net*) wants to help your practice provide optimum care to members and increase HEDIS^{®1}rates. This tip sheet outlines key details of the Use of Imaging Studies for Low Back Pain (LBP) HEDIS measure, its codes and documentation guidelines.

Measure ²	The percentage of memb diagnosis of low back pai study (plain X-ray, MRI, C diagnosis. This measure is reported indicates appropriate tre proportion for whom ima		
Eligible population ²	Age	18 to 75 years as of December 31 of the measurement year. The rate is reported in two age stratifications (18-64 and 65-75) and as a total rate.	
	Intake period	January 1 through December 3 of the measurement year. The intake period is used to identify the first eligible encounter with a principal diagnosis of low back pain.	
	Episode date (anchor date)	The earliest date of service for an eligible encounter during the intake period with a principal diagnosis of low back pain.	
	Product lines	Commercial, Medi-Cal, Medicare	_
• • • • • • • • • • • • •		• • • • • • • • • • • • • • • • • • • •	
Best practice tips	 Educate patients about the second s	about the reason imaging tests are not warra out ways to treat symptoms and prevent reinj ing heavy objects, twisting and bending. Dain relievers. It common low back pain. Irage light walking and complementary care t actic care, acupuncture and/or refer to physic	iury: cherapies such

For member visits where the following events are in evidence, the member would be excluded from the measure. HEDIS exclusions to the LBP measure are shown in the chart below.

	When	Event
	At any time in the patient's history through 28 days after the episode date.	 History of HIV History of cancer Lumbar surgery Osteoporosis therapy or a dispensed prescription to treat osteoporosis Spondylopathy
	Any time during the measurement year.	DeathReceiving palliative careUsing hospice services
Exclusions ²	Within the last 12 months prior to the episode date through 28 days after the episode date.	 History of intravenous drug abuse Major organ transplant Neurologic impairment Spinal infection
	During 90 days prior to the episode date through 28 days after the episode date.	Fragility fractureRecent trauma
	For 90 consecutive days any time during the 366 days prior to and including the episode date.	 Prolonged use of corticosteroids. Corticosteroid medication list:² Hydrocortisone Cortisone Prednisone Prednisolone Methylprednisolone Triamcinolone Dexamethasone Betamethasone/betamethasone acetate
	As of December 31 of the measurement year.	• Ages 66 and older with frailty <u>and</u> advanced illness

For more information

- Refer to the latest NCQA Quality Rating System (QRS) HEDIS Value Set Directory³ for a complete list of diagnosis codes for low back pain.
- Visit the National Institutes of Health Value Set Authority Center at vsac.nlm.nih.gov/welcome or scan this QR code:



¹Healthcare Effectiveness Data and Information Set (HEDIS).
 ²National Committee for Quality Assurance (NCQA) HEDIS Measurement Year 2024 Volume 2: Technical Specifications for Health Plans, Washington, D.C., 2023.
 ³NCQA HEDIS Measurement Year 2024 Volume 2 Value Set Directory, Washington, D.C., 2024.

*Health Net of California, Inc., Health Net Community Solutions, Inc. and Health Net Life Insurance Company are subsidiaries of Health Net, LLC and Centene Corporation. Health Net is a registered service mark of Health Net, LLC. All other identified trademarks/service marks remain the property of their respective companies. All rights reserved.