

Clinical Policy: Semaglutide for Weight Loss (Wegovy)

Reference Number: CP.CPA.352

Effective Date: 09.01.21

Last Review Date: 02.24

Line of Business: Commercial

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Semaglutide (Wegovy®) is a glucagon-like peptide-1 (GLP-1) receptor agonist.

FDA Approved Indication(s)

Wegovy is indicated as an adjunct to a reduced-calorie diet and increased physical activity for chronic weight management in:

- Adult patients with an initial body mass index (BMI) of
 - 30 kg/m² or greater (obesity); or
 - 27 kg/m² or greater (overweight) in the presence of at least one weight-related comorbid condition (e.g., hypertension, type 2 diabetes mellitus, or dyslipidemia);
- Pediatric patients aged 12 years and older with an initial BMI at the 95th percentile or greater standardized for age and sex (obesity).

Limitation(s) of use:

- Wegovy contains semaglutide and should not be coadministered with other semaglutide-containing products or with any other GLP-1 receptor agonist.
- The safety and effectiveness of Wegovy in combination with other products intended for weight loss, including prescription drugs, over-the-counter drugs, and herbal preparations, have not been established.
- Wegovy has not been studied in patients with a history of pancreatitis.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation® that Wegovy is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Weight Management (must meet all):

1. Member meets one of the following (a, b, or c):
 - a. BMI \geq 30 kg/m²;
 - b. BMI \geq 27 kg/m² with at least one indicator of increased cardiovascular risk (e.g., coronary artery/heart disease, hypertension, dyslipidemia, diabetes, elevated waist circumference) or other obesity-related medical condition (e.g., sleep apnea);
 - c. If age is between 12 and 17 years: BMI \geq 95th percentile standardized for age and sex (*see Appendix D*);

CLINICAL POLICY

Semaglutide for Weight Loss

2. Age \geq 12 years;
3. Wegovy is not prescribed concurrently with other semaglutide-containing products or any other GLP-1 receptor agonist(s);
4. Documentation supports member's participation in a Health Net approved weight loss program (e.g., Weight Watchers) or other weight loss programs recommended by the prescriber that involves a reduced calorie diet, increased physical activity, and behavioral modification that meets both of the following (a and b):
 - a. Been actively enrolled in a Health Net approved weight loss program (e.g., Weight Watchers) or other weight loss programs recommended by the prescriber for at least 6 months;
 - b. Will continue to be actively enrolled in a weight loss program while concomitantly prescribed Wegovy;
5. Dose does not exceed the following:
 - a. Week 1 through 4: 0.25 mg once weekly;
 - b. Week 5 through 8: 0.5 mg once weekly;
 - c. Week 9 through 12: 1 mg once weekly;
 - d. Week 13 through 16: 1.7 mg once weekly.

Approval duration: 16 weeks

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial.

II. Continued Therapy

A. Weight Management (must meet all):

1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);

CLINICAL POLICY

Semaglutide for Weight Loss

2. Member is responding positively to therapy as evidenced by one of the following (a or b):
 - a. If this is the first renewal request, member has lost $\geq 5\%$ of baseline body weight (adults) or baseline BMI (pediatrics);
 - b. If this is a second or subsequent renewal request, member has lost weight and/or maintained weight loss on therapy;
3. Wegovy is not prescribed concurrently with other semaglutide-containing products or any other GLP-1 receptor agonist(s);
4. Documentation that member is actively enrolled in a weight loss program that involves a reduced calorie diet, increased physical activity, and behavioral modification adjunct to therapy;
5. Request meets both of the following (a and b):
 - a. Dose does not exceed 2.4 mg once weekly;
 - b. After the initial dose escalation period (*see Section V*), maintenance dose is ≥ 1.7 mg once weekly.

Approval duration: 6 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off-label use policy – CP.CPA.09 or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

BMI: body mass index

FDA: Food and Drug Administration

GLP-1: glucagon-like peptide-1

CLINICAL POLICY
Semaglutide for Weight Loss

Appendix B: Therapeutic Alternatives
Not applicable

Appendix C: Contraindications / Boxed Warnings

- Contraindication(s): personal or family history of medullary thyroid carcinoma (MTC) or with multiple endocrine neoplasia syndrome type 2 (MEN 2), known hypersensitivity reaction to semaglutide or to any of the excipients in Wegovy
- Boxed warning(s): risk of thyroid C-cell tumors

Appendix D: General Information

- BMI = 703 x [weight (lbs)/height (inches)²].
- Examples of coronary artery/heart disease include coronary artery bypass graft, angina, and history of myocardial infarction or stroke.
- The Endocrine Society practice guideline on pharmacological management of obesity states that a weight loss < 5% after 3 months of therapy indicates the weight loss medication is ineffective. In such cases, the Endocrine Society recommends that the medication be discontinued and alternative medications be considered.
- BMI cut-offs (95th percentile) for obesity by age and sex for pediatric patients aged ≥ 12 years:

Age (in years)	95 th Percentile BMI Value	
	Male	Female
12	24.2	25.2
12.5	24.7	25.7
13	25.1	26.3
13.5	25.6	26.8
14	26.0	27.2
14.5	26.4	27.7
15	26.8	28.1
15.5	27.2	28.5
16	27.5	28.9
16.5	27.9	29.3
17	28.2	29.6
17.5	28.6	30.0

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Weight management	<p><u>Adults</u> SC once weekly following dose escalation schedule:</p> <ul style="list-style-type: none"> • Week 1 through 4: 0.25 mg • Week 5 through 8: 0.5 mg • Week 9 through 12: 1 mg • Week 13 through 16: 1.7 mg • Week 17 and onward*: 1.7 mg or 2.4 mg 	2.4 mg/week

CLINICAL POLICY
Semaglutide for Weight Loss

Indication	Dosing Regimen	Maximum Dose
	<p>If patients do not tolerate a dose during dose escalation, consider delaying dose escalation for 4 weeks.</p> <p>The maintenance dosage in adults is either 2.4 mg (recommended) or 1.7 mg once weekly.</p> <p><u>Pediatric patients aged ≥ 12 years old</u> SC once weekly following dose escalation schedule:</p> <ul style="list-style-type: none"> • Week 1 through 4: 0.25 mg • Week 5 through 8: 0.5 mg • Week 9 through 12: 1 mg • Week 13 through 16: 1.7 mg • Week 17 and onward*: 2.4 mg <p>If patients do not tolerate a dose during dose escalation, consider delaying dose escalation for 4 weeks.</p> <p>If patients do not tolerate the 2.4 mg once-weekly maintenance dose, the maintenance dose may be reduced to 1.7 mg once weekly. Discontinue Wegovy if the patient cannot tolerate the 1.7 mg dose.</p> <p><i>* 0.25 mg, 0.5 mg, and 1 mg once-weekly dosages are initiation and escalation dosages and are not approved as maintenance dosages for chronic weight management</i></p>	

VI. Product Availability

Pre-filled, single-dose pens: 0.25 mg, 0.5 mg, 1 mg, 1.7 mg, 2.4 mg

VII. References

1. Wegovy Prescribing Information. Plainsboro, NJ: Novo Nordisk Inc.; July 2023. Available at: www.wegovy.com. Accessed September 5, 2023.
2. Hampl SE, Hassink SG, Skinner AC, et al. Clinical practice guideline for the evaluation and treatment of children and adolescents with obesity [published online ahead of print, 2023 Jan 9]. *Pediatrics*. 2023; e2022060640.
3. Weghuber D, Barrett T, Barrientos-Pérez M, et al. Once-weekly semaglutide in adolescents with obesity. *N Engl J Med*. 2022;387(24):2245-2257.
4. Jensen MD, Ryan DH, Apovian CM, et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *Circulation*. 2014; 129 (suppl 2): S102–S138.

CLINICAL POLICY
Semaglutide for Weight Loss

5. Apovian CM, Aronne LJ, Bessesen DH, et al. Pharmacological management of obesity: an Endocrine Society clinical practice guideline. J Clin Endocrinol Metab. 2015; 100(2): 42-362.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
C9399	Unclassified drugs or biologicals
J3490	Unclassified drugs

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created.	06.15.21	08.21
3Q 2022 annual review: no significant changes; references reviewed and updated.	03.30.22	08.22
Template changes applied to other diagnoses/indications and continued therapy section.	09.29.22	
RT4: updated indication with pediatric expansion of age ≥ 12 years; removed continued therapy criterion of BMI ≥ 25 kg/m ² ; specified continuation therapy positive response criterion of $\geq 5\%$ loss of baseline body weight for adults and BMI for pediatric members.	02.14.23	05.23
3Q 2023 annual review: no significant changes; added HCPCS code; references reviewed and updated.	04.25.23	08.23
Per updated prescribing information, updated dosing to allow and require ≥ 1.7 mg once weekly maintenance dose.	09.05.23	
For documentation of weight loss program, added members has been actively enrolled for at least 6 months, added a weight loss program that also involves behavioral modification, clarified weight loss program to be either a Health Net approved weight loss program or a weight loss program recommended by the prescriber.	12.12.23	02.24

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in

CLINICAL POLICY

Semaglutide for Weight Loss

developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions. Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2021 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.